

Table 33. Estimated Number of Persons Living with AIDS¹ by Sex and Exposure Category, 1993–98

	Exposure Category							Total	Percent drug-related ³
	Men who have sex with men (MSM)	Injecting drug use (IDU)	MSM and IDU	Hemophilia/coagulation disorder	Heterosexual contact	Receipt of blood transfusion ²	Risk not reported or identified		
Male adult/adolescent									
1993	86,720	34,465	13,645	1,618	6,066	918	1,015	144,467	33.3%
1994	95,171	40,153	14,635	1,698	7,690	947	969	161,463	33.9%
1995	102,717	44,589	15,369	1,728	9,771	1,008	969	175,151	34.2%
1996	111,076	49,074	16,034	1,742	12,219	1,083	1,011	192,239	33.8%
1997	122,874	54,249	17,203	1,788	15,011	1,189	1,064	213,378	33.4%
1998	133,840	58,843	18,181	1,838	17,765	1,313	1,133	232,913	33.0%
Female adult/adolescent									
1993	N/A	13,793	N/A	91	11,829	766	383	26,862	51.3%
1994	N/A	16,175	N/A	108	16,175	854	390	32,667	49.5%
1995	N/A	18,294	N/A	137	18,524	908	392	38,255	47.8%
1996	N/A	20,285	N/A	167	22,588	1,006	429	44,475	45.6%
1997	N/A	22,586	N/A	207	27,017	1,120	477	51,407	43.9%
1998	N/A	24,500	N/A	240	31,336	1,243	519	57,838	42.3%

N/A Not applicable.

¹Excludes pediatric AIDS cases (<13 years old). These numbers do not represent actual cases of persons living with AIDS. Rather, these numbers are point estimates of persons living with AIDS derived by subtracting the estimated cumulative number of deaths in persons with AIDS from the estimated cumulative number of persons with AIDS. Estimated AIDS cases are adjusted for reporting delays and for redistribution of cases initially reported with no identified risk, but not for incomplete reporting. Annual estimates are through the most recent year for which reliable estimates are available.

²Includes receipt of blood components or tissue.

³Proportion includes injection drug users and MSM who are injection drug users.

Source: Centers for Disease Control and Prevention. *HIV/AIDS Surveillance Report, 1999*; 11 (No. 2), Table 28.